

Altered brain connectivity in dysmenorrhea: pain modulation and the motor cortex

Jason J. Kutch^a, Frank F. Tu^{b,c}

Primarily dysmenorrhea, menstrual pain without an identified cause, is a common and often debilitating problem. Fifteen to twenty percentage of women fail to respond to conservative treatment and experience significant reduction in quality of life.⁵ An important consideration, that we and others have proposed, is that these treatment failures may reflect a completely different phenotype, where the primary disorder is aberrant central processing of pain rather than solely peripherally based uterine inflammation.⁴ Consistent with this idea, high rates of comorbidity have been identified between dysmenorrhea and other idiopathic pain disorders, notably irritable bowel syndrome and painful bladder syndrome.¹⁸ The epidemiological overlap with these visceral pain conditions suggests that sensitization in the central nervous system may also play a role in menstrual uterine pain. Correspondingly, there is a growing body of evidence showing that some dysmenorrhea sufferers exhibit widespread reductions in both somatic^{2,6} and visceral¹⁶ hyperalgesia, suggesting fundamental changes in central pain modulatory systems.

In this issue, Wei et al.¹⁷ tested the hypothesis that brain networks, known to modulate pain, might display markers of dysfunction in women with primary dysmenorrhea. They performed resting state functional magnetic resonance imaging of the brain in women both with and without primary dysmenorrhea at 2 time points: during the menstrual phase and during the periovulatory phase. Resting state functional magnetic resonance imaging is an established approach to estimate strength of interaction (functional connectivity) among brain regions. As in previous studies, by examining women at these 2 time points, the authors were able to disambiguate functional connectivity changes related to the active experience of pain during the menstrual phase from functional connectivity changes more intrinsic to the individual during the periovulatory phase. Due to the wealth of literature showing a role for periaqueductal gray (PAG) in descending pain modulation, the authors hypothesized that the functional connectivity of PAG might distinguish women with and without primary dysmenorrhea. During the menstrual phase, women with primary dysmenorrhea had increased functional connectivity, compared with healthy controls, between the PAG and medial sensorimotor areas near the midline of the brain (stronger changes are reported in motor compared with

sensory areas). However, during the periovulatory phase, women with primary dysmenorrhea had decreased functional connectivity with the “default mode network,” a group of brain regions known to be important in healthy brain function.

A particularly exciting feature of the report of Wei et al. is potentially convergent brain imaging findings across chronic visceral pain disorders. Recent studies have reported functional connectivity disturbances in medial brain motor areas in women with interstitial cystitis/painful bladder syndrome,⁹ in men with chronic prostatitis/chronic pelvic pain syndrome,¹⁰ and also in women with localized provoked vulvodynia.⁷ Recent anatomical connectivity studies in rats using retrograde transneuronal labelling studies suggest that visceral structures may be more strongly represented in medial brain motor areas than any other cortical area; also, the visceral representation in motor cortex is aligned with the more medial somatic motor representation of trunk and lower limb muscles.¹² In humans, muscles, such as the pelvic floor,¹ abdominals,¹⁵ and toes,¹⁴ are represented in more medial areas of motor cortex, and dysfunction of these regions has been suggested in chronic pain.⁸ These data collectively suggest that medial brain motor areas may have important but underrecognized modulatory influence on both somatic and visceral processes.

The specific relationship between PAG and the motor cortex in dysmenorrhea is also supported by recent reviews of therapeutic transcranial magnetic stimulation. Motor cortex is the only brain region for which strong evidence exists for the analgesic effects of transcranial magnetic stimulation, possibly through the indirect (multisynaptic) interaction of motor cortex with the PAG.¹¹ The findings of Wei et al. in this issue might suggest that the pain of primary dysmenorrhea during the menstrual phase could relate to a dysfunctional pain modulatory system involving the motor cortex and PAG. If the importance of connections between motor cortex and the viscera shown in animal models can be repeatedly demonstrated in humans, it is possible that motor cortical dysfunction in visceral pain conditions could result in a wide array of effects on somatic motor activity, visceral motor activity, and descending pain modulation.

The study of Wei et al. in this issue opens interesting questions. It remains unclear if the PAG functional connectivity changes during the menstrual phase are more or less important in mediating symptoms compared with the changes during the periovulatory phase. Functional connectivity changes within the default mode network during the periovulatory phase are intriguing given the extensive previous literature on the default mode network in chronic pain^{3,13} and that these changes exist even when an individual is not experiencing the active pain of the menstrual phase (ie, may represent a trait of the individual). It is possible that PAG functional connectivity changes to the default mode network may be an ongoing representation of cumulative

Sponsorships or competing interests that may be relevant to content are disclosed at the end of this article.

^a Division of Biokinesiology and Physical Therapy, University of Southern California, Los Angeles, CA, USA, ^b Department of Obstetrics and Gynecology, NorthShore University HealthSystem, Evanston, IL, USA, ^c The Pritzker School of Medicine, University of Chicago, Chicago, IL, USA

PAIN 157 (2016) 5–6

© 2015 International Association for the Study of Pain

<http://dx.doi.org/10.1097/j.pain.0000000000000364>

pain experienced during the menstrual phase, pain which is actually generated by functional connectivity dysfunction between medial brain motor regions and the PAG. Future longitudinal studies with repeated neuroimaging during both the menstrual and periovulatory phases, and associated symptoms, will be necessary to tease out the relative importance of PAG functional connectivity to the default mode network compared with medial brain motor areas. The work of Wei et al. is nonetheless very important to stimulate future research in motor cortical involvement in pain processes.

Conflict of interest statement

The authors have no conflicts of interest to declare.

Acknowledgements

J. J. Kutch acknowledges support from Grant U01DK082370 and F. F. Tu acknowledges support from Grant 1R01DK100368, both from the National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, in the writing of this article. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Article history:

Received 4 September 2015

Accepted 8 September 2015

Available online 21 September 2015

References

- [1] Asavasopon S, Rana M, Kirages D, Yani M, Fisher B, Hwang D, Lohman E, Berk L, Kutch J. Cortical Activation Associated with Muscle Synergies of the Human Male Pelvic Floor. *J Neurosci* 2014;34:13811–18.
- [2] Bajaj P, Bajaj P, Madsen H, Arendt-Nielsen L. A comparison of modality-specific somatosensory changes during menstruation in dysmenorrheic and nondysmenorrheic women. *Clin J pain* 2002;18:180–90.
- [3] Baliki MN, Geha PY, Apkarian AV, Chialvo DR. Beyond feeling: chronic pain hurts the brain, disrupting the default-mode network dynamics. *J Neurosci* 2008;28:1398–403.
- [4] Bettendorf B, Shay S, Tu F. Dysmenorrhea: contemporary perspectives. *Obstet Gynecol Surv* 2008;63:597–603.
- [5] Dawood MY. Primary dysmenorrhea: advances in pathogenesis and management. *Obstet Gynecol* 2006;108:428–41.
- [6] Giamberardino MA, Berkley KJ, Iezzi S, de Bigontina P, Vecchiet L. Pain threshold variations in somatic wall tissues as a function of menstrual cycle, segmental site and tissue depth in non-dysmenorrheic women, dysmenorrheic women and men. *PAIN* 1997;71:187–97.
- [7] Gupta A, Rapkin AJ, Gill Z, Kilpatrick L, Fling C, Stains J, Masghati S, Tillisch K, Mayer EA, Labus JS. Disease-related differences in resting-state networks: a comparison between localized provoked vulvodynia, irritable bowel syndrome, and healthy control subjects. *PAIN* 2015;156:809–19.
- [8] Hodges PW, Tucker K. Moving differently in pain: a new theory to explain the adaptation to pain. *PAIN* 2011;152(3 suppl):s90–8.
- [9] Kilpatrick L, Kutch J, Tillisch K, Naliboff B, Labus J, Jiang Z, Farmer M, Apkarian A, Mackey S, Martucci K, Clauw D, Harris R, Deutsch G, Ness T, Yang C, Maravilla K, Mullins C, Mayer E. Alterations in Resting State Oscillations and Connectivity in Sensory and Motor Networks in Women with Interstitial Cystitis/Painful Bladder Syndrome. *J Urol* 2014;192:947–55.
- [10] Kutch J, Yani M, Asavasopon S, Kirages D, Rana M, Cosand L, Labus J, Kilpatrick L, Ashe-McNailey C, Farmer M, Johnson K, Ness T, Deutsch G, Harris R, Apkarian A, Clauw D, Mackey S, Mullins C, Mayer E. Altered resting state neuromotor connectivity in men with chronic prostatitis/chronic pelvic pain syndrome: a MAPP Research Network Neuroimaging Study. *Neuroimage Clin* 2015;8:493–502.
- [11] Lefaucheur J-P, André-Obadia N, Antal A, Ayache SS, Baeken C, Benninger DH, Cantello RM, Cincotta M, De Carvalho M, De Ridder D. Evidence-based guidelines on the therapeutic use of repetitive transcranial magnetic stimulation (rTMS). *Clin Neurophysiol* 2014;125:2150–206.
- [12] Levinthal DJ, Strick PL. The motor cortex communicates with the kidney. *J Neurosci* 2012;32:6726–31.
- [13] Napadow V, LaCount L, Park K, As-Sanie S, Clauw DJ, Harris RE. Intrinsic brain connectivity in fibromyalgia is associated with chronic pain intensity. *Arthritis Rheum* 2010;62:2545–55.
- [14] Schrum A, Wolff S, van der Horst C, Kuhtz-Buschbeck JP. Motor Cortical Representation of the Pelvic Floor Muscles. *J Urol* 2011;186:185–90.
- [15] Tsao H, Galea M, Hodges P. Reorganization of the motor cortex is associated with postural control deficits in recurrent low back pain. *Brain* 2008;131:2161–71.
- [16] Tu FF, Epstein AE, Pozolo KE, Sexton DL, Melnyk AI, Hellman KM. A noninvasive bladder sensory test supports a role for dysmenorrhea increasing bladder noxious mechanosensitivity. *Clin J Pain* 2013;29:883–90.
- [17] Wei SY, Chao HT, Tu CH, Li WC, Low I, Chuang CY, Chen LF, Hsieh JC. Changes in functional connectivity of pain modulatory systems in women with primary dysmenorrhea. *PAIN* 2015;157:90–100.
- [18] Zondervan KT, Yudkin PL, Vessey MP, Jenkinson CP, Dawes MG, Barlow DH, Kennedy SH. The community prevalence of chronic pelvic pain in women and associated illness behaviour. *Br J Gen Pract* 2001;51:541–7.